

Soybean - Optimum Planting Conditions and Seed Placement

- For optimal emergence and stand establishment, soybean seed should be planted when soil temperatures are warm enough for germination.
- Planting depth, row spacing, and planting population are important for stand establishment and plant development.
- Farmers can determine when and how to plant their crop; therefore, controlling many management factors that may improve soybean yield potential.

Optimal Soil Temperatures

Soybean seeds can germinate when soil temperatures reach at least 54° F, but a temperature of 60° F is optimal.¹ When soil temperatures are too cool, seeds stay dormant and become more vulnerable to diseases, insects, and animal predators. Soybean seed should be planted when soils have had time to warm and within the target dates for a region. Planting into cold and/ or wet soils can lead to numerous problems.

Saturated and Flooded Soils

Soybean seed should only be planted when adequate soil moisture is available and high soil oxygen concentrations are favorable for germination and plant growth. In some southern and western regions planting is often driven by soil moisture content. Saturated soils, which may include flooded or ponded soils, can have a negative impact on emergence. The main side effects include plant growth restriction and decreased oxygen availability to the plant.

Saturated soils can inhibit root growth, leaf area expansion, and the photosynthetic process. When soils become saturated, young plants may develop yellow leaves due to slowing of photosynthesis and plant growth. A prolonged period of saturated soil can reduce germination and emergence due to lack of oxygen. In addition, portions of roots may die as a result of no oxygen. However, there is still a chance for survival unless the growing point is damaged.

The longer an area remains saturated, the higher the risk of plant death, and warm air temperatures during soil saturation can shorten plant survival.² For soybean, yield losses are not typically noted in fields flooded for 2 days or less. Flooding for 4 days or more can stress the crop, delay plant growth, and cause shorter plants with fewer nodes. Six days of flooding can cause significant soybean

yield loss and flooding for a week or more can result in entire loss of the stand.

Soil Compaction

Soil compaction takes place when soil particles are pressed together resulting in decreased pore space and increased soil density. Compaction can result in yield reductions due to decreases in seed germination, root and plant growth, and nutrient uptake (Figure 1). The majority of soil compaction is created from equipment passes over a field. Research indicates that approximately 80% of soil compaction happens on the first pass, while subsequent passes cause additional, but progressively less, compaction.³ The way to manage compaction is prevention. That means staying out of the field until conditions for field work have improved. If mud sticks to the tires and ruts are deeper than an inch, it is too wet to be in the field.



Figure 1. Effect of compaction on soybean growth and nutrient uptake.

Soybean - Optimum Planting Conditions and Seed Placement

Soil Crusting

As wet soils dry, a crust layer can form on the surface. The crust layer can delay or prevent seedlings from emerging. In addition, soybean hypocotyls can easily be broken when trying to push through the crust. Crusting may be more common in fields with fine textured soils, low organic matter, and little surface residue.⁴ A rotary hoe can break up the crust and aid in seedling emergence. Timing is essential, and breaking the crust as soon as possible is most beneficial to the young plants. Cooler soils allow seedlings infected with disease to survive longer when trying to break through the crust.

Planting

Soybean seed should never be planted too deep. Generally, soybean seed should be planted 1 to 1.5 inches deep and no deeper than 2 inches. Soybean plants emerge as the hypocotyl straightens and carries the cotyledons to the surface. The plant requires a lot of energy to complete this process. Therefore, planting too deep can burn energy that could be used later by the plant. In addition, planting too deep can inhibit emergence in stressful situations, such as soil crusting and compaction.

Summary

Planting when soil temperature and conditions are favorable, at the proper depth, seeding rate, and row configuration is very important to give the crop the best chance for emergence and stand establishment. Waiting for warm soil temperatures and conditions may help seedlings avoid diseases, reduced oxygen situations, and other plant growth restrictions. Although it is important to plant within the acceptable planting window, rushing to plant in cold, wet conditions can lead to yield-reducing problems later in the season.

Sources:

¹Louisiana State University. 2003. Crop profile for soybeans in Louisiana. www.ipmcenters.org (verified 3/4/14).

²Pedersen, P. 2007. Planting and replanting scenarios. Iowa State University Extension. Integrated Crop Management. IC-498(13). www.ipm.iastate.edu (verified 3/4/14).

³Daum, D.R. 1996. Soil compaction and conservation tillage. Penn State Cooperative Extension. <http://extension.psu.edu> (verified 3/4/14).

⁴Al-Kaisi, M. and P. Pedersen. 2007. Wet conditions: Challenges and opportunities. Iowa State University Extension. Integrated Crop Management. IC-498 (9). www.ipm.iastate.edu (verified 3/4/14).

⁵Bogren, R. 2008. Researchers say don't plant soybeans too early. Soybean and Grain Promotion Board Reports. Louisiana State University Ag Center. www.lsuagcenter.com (verified 3/4/14).

⁶Clawson, L. and D. J. Boquet. 2007. Louisiana State University Ag Center. Winter Issue.

For additional agronomic information, please contact your local seed representative.

Individual results may vary, and performance may vary from location to location and from year to year. This result may not be an indicator of results you may obtain as local growing, soil and weather conditions may vary. Growers should evaluate data from multiple locations and years whenever possible. **ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Leaf Design[®] is a registered trademark of Monsanto Company. All other trademarks are the property of their respective owners. ©2015 Monsanto Company. 140216060210 030614CRB.