

General Characteristics

| | |
|--------------------|-----------|
| Winter Survival* | 1.5 |
| Recovery | Very Fast |
| Persistence | Excellent |
| Leaf to Stem Ratio | Excellent |
| Fall Dormancy** | 4 |

*Winter Survival-Lower number - more persistent variety

**Fall Dormancy-Higher number - earlier it starts growing in spring, later it grows in the fall

Agronomic Features

| | |
|--------------------------|-----------|
| Maturity | Med-Early |
| Forage Yield | Very Good |
| Dry Down | Excellent |
| Stress Tolerance | Very Good |
| Traffic Tolerance | Excellent |
| Standability | Excellent |
| Multileaf Expression | Excellent |
| Digestibility/Feed Value | Very Good |

Disease Resistance

| | |
|--------------------------|-------|
| Bacterial Wilt | HR |
| Verticillium Wilt | HR |
| Fusarium Wilt | HR |
| Anthracnose | HR |
| Phytophthora Root Rot | HR |
| Aphanomyces Root Rot | HR |
| Disease Resistance Index | 35/35 |

Recommended Seeding Rates

| | |
|--------------------|-------|
| Alone (pure stand) | 18-22 |
| In a mix (grasses) | 8-16 |



Product Highlights

- Rated for traffic tolerance
- Wide area of adaptation and soil types
- High leaf to stem ratio
- Branch rooted variety

Pest Resistance Traits

